



TITLE:

Effective use of high CO efflux at the soil surface in a tropical understory plant.

AUTHOR(S):

Ishida, Atsushi; Nakano, Takashi; Adachi, Minaco;
Yoshimura, Kenichi; Osada, Noriyuki; Ladpala, Phnumard;
Diloksumpun, Sapit; Puangchit, Ladawan; Yoshimura, Jin

CITATION:

Ishida, Atsushi ...[et al]. Effective use of high CO efflux at the soil surface in a tropical understory plant.. Scientific reports 2015, 5: 8991.

ISSUE DATE:

2015-03-11

URL:

<http://hdl.handle.net/2433/198826>

RIGHT:

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

OPEN

SUBJECT AREAS:

C₃ PHOTOSYNTHESIS
ECOPHYSIOLOGY

Received
29 September 2014

Accepted
10 February 2015

Published
11 March 2015

Correspondence and
requests for materials
should be addressed to
A.I. (atto@ecology.
kyoto-u.ac.jp)

Effective use of high CO₂ efflux at the soil surface in a tropical understory plant

Atsushi Ishida¹, Takashi Nakano², Minaco Adachi³, Kenichi Yoshimura⁴, Noriyuki Osada⁵,
Phanumard Ladpala⁶, Sapit Diloksumpun⁷, Ladawan Puangchit⁷ & Jin Yoshimura^{8,9,10}

¹Center for Ecological Research, Kyoto University, Shiga 520-2113, Japan, ²Mount Fuji Research Institute, Yamanashi 403-0005, Japan, ³Institute of Industrial Science, The University of Tokyo, Komaba, Tokyo 153-8505, Japan, ⁴Kansai Research Center, Forestry and Forest Products Research Institute, Kyoto 612-0855, Japan, ⁵Tomakomai Experimental Forest, Field Center for Northern Biosphere, Hokkaido University, Tomakomai 053-0035, Japan, ⁶National Park, Wildlife and Plant Conservation Department, Bangkok 10900, Thailand, ⁷Faculty of Forestry, Kasetsart University, Bangkok 10900, Thailand, ⁸Graduate School of Science and Technology, and Department of Mathematical and Systems Engineering, Shizuoka University, Shizuoka 432-8561, Japan, ⁹Department of Environmental Science and Forest Biology, State University of New York College of Environmental Science and Forestry, Syracuse, NY 13210, USA, ¹⁰Marine Biosystems Research Center, Chiba University, Kamogawa, Chiba 299-5502, Japan.

Many terrestrial plants are C₃ plants that evolved in the Mesozoic Era when atmospheric CO₂ concentrations ([CO₂]) were high. Given current conditions, C₃ plants can no longer benefit from high ambient [CO₂]. *Kaempferia marginata* Carey is a unique understory ginger plant in the tropical dry forests of Thailand. The plant has two large flat leaves that spread on the soil surface. We found a large difference in [CO₂] between the partly closed space between the soil surface and the leaves (638 μmol mol⁻¹) and the atmosphere at 20 cm above ground level (412 μmol mol⁻¹). This finding indicates that the plants capture CO₂ efflux from the soil. Almost all of the stomata are located on the abaxial leaf surface. When ambient air [CO₂] was experimentally increased from 400 to 600 μmol mol⁻¹, net photosynthetic rates increased by 45 to 48% under near light-saturated conditions. No significant increase was observed under low light conditions. These data demonstrate that the unique leaf structure enhances carbon gain by trapping soil CO₂ efflux at stomatal sites under relatively high light conditions, suggesting that ambient air [CO₂] can serve as an important selective agent for terrestrial C₃ plants.

The geological record indicates that the C₃ land plants originated during the middle to late Ordovician period (450 to 440 million years ago) when atmospheric CO₂ concentrations ([CO₂]) were still very high (approximately 4% compared with 0.039% at present) and O₂ concentrations ([O₂]) in air were low (approximately 15% compared with 21% at present)^{1,2}. Although the down-regulation of Rubisco (ribulose-1,5-bisphosphate carboxylase/oxygenase) under high [CO₂] is a well-known phenomenon³, high [CO₂] and low [O₂] in the ancient air would have contributed to an increase in carbon assimilation rates (A) due to the kinetics of Rubisco. A meta-analysis of FACE (free-air CO₂ enrichment) experiments revealed that the average maximum carboxylation rates under doubled [CO₂] were -17% in C₃ crops and -4% in C₃ trees due to dawn-regulation. On average, the increase in light-saturated net photosynthesis under doubled [CO₂] was 13% in C₃ crops and 47% in C₃ trees³. This finding may indicate that C₃ plants in the past exhibited increased carbon (C) gain and that more extensive C cycling occurred in forest ecosystems compared with the present era. During the Cenozoic era, atmospheric O₂ concentrations increased and atmospheric [CO₂] became largely depleted, with record minimum [CO₂] during the Oligocene/Miocene epoch (24 million years ago)⁴. Since the advent of the Industrial Revolution, atmospheric [CO₂] has increased rather rapidly due to the modernization of human society and increasing reliance on coal and oil burning. In the photosynthetic CO₂-response curves of C₃ plants, the transition of the limitation from ribulose-1,5-bisphosphate (RuBP) carboxylation limitation to RuBP regeneration limitation is typically observed between ambient and doubled ambient [CO₂]⁵. Thus, C₃ plants are constrained by the carboxylation limit of RuBP in the present-day air [CO₂]. In contrast, photosynthesis in C₄ plants is not limited by low air [CO₂]⁶ because these plants possess the appropriate enzyme (PEP carboxylase) and the specific anatomy in bundle sheath cells required to increase the CO₂ partial pressure around Rubisco sites⁷. C₄ plants have evolved to improve plant carbon and water relations simultaneously during photosynthesis and to cope with declining atmospheric [CO₂] and increasing water demand^{4,8,9}. However, C₃ plants have not evolved carbon-concentrating mechanisms in their physiology and anatomy.

Even in present-day ecosystems, sites with high air $[\text{CO}_2]$, such as forest floors¹⁰ and volcanic vents¹¹ are observed. The high $[\text{CO}_2]$ found on forest floors originates from the respiration of soil organisms and plant-root systems. Attention has been focused on the large contributions of sunflecks or sun patches to net C assimilation rates (A) in forest understory plants, indicating strong light limitation^{12,13}. However, the potential effects of rising $[\text{CO}_2]$ on A in understory plants have rarely been evaluated. High $[\text{CO}_2]$ should contribute to the survival of understory plants that experience reduced photosynthetic rates due to water stress¹⁴. The stable carbon isotope ratios of understory plants indicate that these plants re-fix the efflux C in tropical¹⁵ and cool-temperate forests¹⁶.

High $[\text{CO}_2]$ that originates from the soil surface dissipates rapidly due to diffusion and mass flow caused by wind. Although wind velocity is reduced near the understory, an extremely gentle breeze is sufficient to diffuse CO_2 from the soil surface¹⁷. Therefore, for understory plants to effectively use this high soil-efflux $[\text{CO}_2]$, they must trap CO_2 near the soil surface. In the present study, we report the discovery of an understory ginger plant, *Kaempferia marginata* Carey (Zingiberaceae), which effectively traps soil-efflux CO_2 in the closed space between the soil surface and its leaves. This plant enhances photosynthesis by 45 to 48% under relatively high light conditions. It is a drought-deciduous, perennial herb found in tropical dry forests in Southeast Asia. Based on measurements of ambient air $[\text{CO}_2]$, photosynthetic capacity, and the stable carbon isotope ratios in the lamina, we demonstrate that this ginger plant makes effective use of high $[\text{CO}_2]$ on the forest floor.

Results

The ginger plant has a unique leaf structure; the individual plant has two flat leaves that spread on the soil surface, and the leaf edges are often curled downward to capture the air under its leaf blades (Fig. 1). The root system is small, indicating that this plant has a poor water uptake capacity. The uppermost height of a single leaf blade is only 24 mm above the ground surface on average and defines a relatively closed space between the leaf blade and the soil surface (Table S1). The stomatal densities were 1.6 mm^{-2} and 20.9 mm^{-2} on the adaxial




Figure 1 | An understory ginger plant, *Kaempferia marginata* Carey, with a unique leaf structure in a tropical forest in Southeast Asia. (a) field-grown plants, (b) A plant removed from the soil; two large leaves and a poor root system are evident.

and abaxial leaf surfaces, respectively, indicating that approximately all stomata face the soil surface. The distributions of leaf sizes and leaf morphologies indicate that as the leaf size increases with time, the leaf shape gradually becomes rounder (Fig. S1), contributing to an increase in the efficiency of trapping CO_2 efflux from the soil surface.

On a sunny day during the rainy season, the average daily $[\text{CO}_2]$ was $412 \mu\text{mol mol}^{-1}$ in the open air at 20 cm above the ground and $638 \mu\text{mol mol}^{-1}$ in the space between the leaves and soil surface (Fig. 2). The maximum $[\text{CO}_2]$ observed in the air space was greater than $1000 \mu\text{mol mol}^{-1}$. Nevertheless, $[\text{CO}_2]$ in the space largely fluctuated with temporal variations in wind velocity. The values (mean \pm SD) of the stable carbon isotope ratios ($\delta^{13}\text{C}$) in the lamina were $-34.9 \pm 1.5 \text{ ‰}$ in the ginger plants and $-29.1 \pm 1.5 \text{ ‰}$ in the upper canopy leaves of woody plants in the dry evergreen forest (our unpublished data on woody plants). The low $\delta^{13}\text{C}$ value in the ginger plants indicates high internal $[\text{CO}_2]$ in the leaves during the day.

When the ambient-air $[\text{CO}_2]$ was artificially increased from 400 to $600 \mu\text{mol mol}^{-1}$, the A under near-light saturated conditions ($800 \mu\text{mol m}^{-2} \text{ s}^{-1}$ PPF: photosynthetic photon flux) increased from 5.8 to $8.2 \mu\text{mol m}^{-2} \text{ s}^{-1}$, a 45% increase (Fig. 3A). In contrast, under low light conditions (less than $70 \mu\text{mol m}^{-2} \text{ s}^{-1}$ PPF), no significant increase was detected in A after elevating $[\text{CO}_2]$ from 400 to $600 \mu\text{mol mol}^{-1}$. We also measured ambient-air CO_2 response curves under 500 and $40 \mu\text{mol m}^{-2} \text{ s}^{-1}$ PPFs. Both RuBP carboxylation and RuBP regeneration rates were reduced by the low PPF (Fig. 3B). When the ambient-air $[\text{CO}_2]$ was increased from 400 to $600 \mu\text{mol mol}^{-1}$, A increased by 48% under relatively strong sunlight ($500 \mu\text{mol m}^{-2} \text{ s}^{-1}$ PPF) and by 36% under reduced light ($40 \mu\text{mol m}^{-2} \text{ s}^{-1}$ PPF) conditions. The data indicate that a significant increase in A in response to elevated $[\text{CO}_2]$ was more pronounced under sunlit conditions compared with shaded conditions. Sunflecks must thus cooperate with rising $[\text{CO}_2]$ for enhancing of A ^{12,13}.

Discussion

The data presented here indicate that the unique leaf structure of ginger plant enhances C fixation under high light conditions by effectively trapping high $[\text{CO}_2]$ efflux in the relatively closed space between their leaves and the soil surfaces. In tropical forests, high termite activity at ground level prevents fallen leaves from covering the leaf surface of the ginger plants (Fig. 1A); the leaf litter layer typically remains fairly thin and does not persist for a long period of time. This may be a factor in explaining why the ginger plant has evolved to capture CO_2 efflux from soil respiration in tropical forests.

Another unique morphological characteristic of the ginger plant is the small root system (Fig. 1B). Large non-photosynthetic organs are found to have large respiration requirements^{18,19}. However, its small root system, the ginger plant has a very low CO_2 compensation point at the whole plant level, similar to leafy plants¹⁴. Because of the small root system, the ginger plant can only grow during the favorable rainy season as an ephemeral plant. Another advantage is the high




Figure 2 | Diurnal time variations in air CO_2 concentration at 20 cm above the ground (open circles) and in the air space between the leaf blade and soil surface (blue circles).




Figure 3 | Photosynthetic responses under high CO_2 concentrations and high light conditions. (a) Photosynthetic light response curves (PPF- A curves) at 600 $\mu\text{mol mol}^{-1}$ CO_2 (closed circles) and 400 $\mu\text{mol mol}^{-1}$ CO_2 (open circles), where PPF represents the photosynthetic photon flux at the leaf surface and A is net C assimilation rate. Bars indicate 1 SD, (b) Photosynthetic ambient-air CO_2 response curves (C_a - A curves) in a leaf blade at 500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PPF (closed circles) and 40 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PPF (open circles), where C_a represents CO_2 concentration in outlet gas stream in the LI-6400, i.e., ambient air CO_2 concentration.

soil respiration during rainy seasons. In tropical dry forests in Thailand, where the ginger plant is a native species, the soil respiration rates become double during the rainy seasons²⁰. The mean soil respiration rate is approximately 7.67 $\mu\text{mol m}^{-2} \text{s}^{-1}$ in the rainy season and approximately 3.63 $\mu\text{mol m}^{-2} \text{s}^{-1}$ in the dry season.

A relatively high irradiance is required to effectively enhance A under elevated $[\text{CO}_2]$ (Fig. 3B); light levels greater than approximately 6.4% of full sunlight appear to be required to maintain a population of the ginger plant (see Environmental description in Supplementary information). Under sunlit conditions, the risk of photoinhibition increases even in tropical climates, particularly in shaded plants at relatively high temperatures^{21,22}. However, in the ginger plant, xanthophyll-cycle dependent non-photochemical quenching (NPQ) appears to prevent chronic photoinhibition (Fig. S3). This unique adaptation to specific microhabitats is reflected by the plant distribution. In the tropical dry forests, the ginger plant is primarily located in the drought-deciduous forests with sparse tree cover and lightly shaded forest floors. In contrast, the ginger plant is exclusively located on the edges of dry evergreen forests with closed canopies.

The discovery of the morphological adaptation of the ginger plant is the first demonstration of the effective use of high CO_2 efflux from soil in understory C_3 plants. Their unique structure of this plant is characterized by large, flat leaves, thus earning the nickname “terrestrial water lily”. The shape delimits the space between the leaves and the ground surface (Fig. 1A). Plants with such an ideal leaf structure are rare even in the tropics. We suggest that the C_3 ginger plant evolved to cope with low atmospheric CO_2 by morphologically trapping high CO_2 efflux from the soil, whereas C_4 plants did so by physiologically concentrating CO_2 within the plant body. In adult trees of certain woody plants, the respiration rates per unit stem surface at breast height ranges from 1.2 $\mu\text{mol m}^{-2} \text{s}^{-1}$ to 3.5 $\mu\text{mol m}^{-2} \text{s}^{-1}$ ^{23,24,25}. The CO_2 efflux from the stem surface is due to numerous parenchyma cells located within stems¹⁸, carbon transport in phloem from the leaves to the roots²⁶, and CO_2 up-flow from root systems due to transpiration-driven sap flow^{20,27,28,29}. Although the stem respiration rates of large trees are reduced compared with than the soil respiration rates during the rainy season, stem respiration may be valuable as a CO_2 source for living plants. Therefore, a mechanism similar to that of the ginger plant may be identified among the lichens, mosses, ferns, orchids, and vines growing not only on the ground but also on the trunks of large trees. We hypothesize that the combination of a closed air space and relatively high sunlight is required to exploit extremely high efflux CO_2 .

The pulse-labeling method has been used to determine the time lag from CO_2 efflux from soil to leaf C assimilation³⁰. The time lag ranges from 12.5 ± 7.5 (mean \pm SD) h in grasses to 4 to 5 days in trees. Although the data indicate that interactions between the soil and plants in the C cycles within a single ecosystem exist, most CO_2 that originates from the soil will have dissipated from the ecosystem by diffusion during this time period. The low $\delta^{13}\text{C}$ values of ginger plants indicate that they were exposed to high $[\text{CO}_2]$ and used large amounts of C emitted from the soil. Nevertheless, shady conditions increase internal $[\text{CO}_2]$ in leaves due to the reduced A , consequently decreasing the $\delta^{13}\text{C}$ values in laminae³¹. Therefore, we cannot use $\delta^{13}\text{C}$ values to distinguish between the two potential sources of the effects, shade and high ambient air $[\text{CO}_2]$. Overall, we can conclude that root and microbial-derived CO_2 are major contributors to carbon assimilation in this ginger plant.

Methods

The study was conducted in July 2008 in a dry evergreen forest in Thailand ($14^\circ 29' \text{N}$, $101^\circ 55' \text{E}$, 563 m ASL) approximately 180 km northeast of Bangkok during the middle of the rainy season³². We selected a population of ginger plants found roadside in a forest with a dense canopy. During three successive days, the diurnal time courses of PPF, ambient air temperatures and relative humidity in air were measured near the center of the plant population (data shown in Fig. S2). On a relatively sunny day, the diurnal time courses of leaf gas exchange and chlorophyll fluorescence were measured from predawn to dusk using an open, portable measurement system (LI-6400, LI-COR, Lincoln, NE) and a chlorophyll fluorescence meter (Mini-PAM, Walz, Effeltrich, Germany), respectively. These measurements were conducted in eight individual plants with relatively large leaves.

While measuring diurnal leaf gas exchange, the diurnal variations in ambient air $[\text{CO}_2]$ were simultaneously measured with thin-film capacitance CO_2 sensors (GM70, Vaisala, Helsinki, Finland) without tube-absorbing air. The CO_2 sensors were set at two heights: 1) 20 cm above the ground and 2) in the air space between the leaf blade and the ground surface in an individual plant with a relatively large leaf area. The diameter of the CO_2 sensor probe was 18.5 mm, and the leaf diameter was greater than 100 mm. Because of without tube-absorbing and given a large leaf, $[\text{CO}_2]$ in the air space below the leaf could be directly measured (Fig. S4); it is possible that we did not completely avoid air leaks along the side of the probe, possibly resulting in an underestimation of $[\text{CO}_2]$.

In the following days, to evaluate the interactive effects of light intensity and $[\text{CO}_2]$ on A , we measured photosynthetic light responses (PPF- A curve) under different ambient air $[\text{CO}_2]$ levels and photosynthetic ambient air CO_2 responses (C_a - A curve) under different light levels during the daylight hours (C_a refers to ambient air $[\text{CO}_2]$). To evaluate the average internal $[\text{CO}_2]$ in leaves over a long time period, carbon isotope ratios in the eight laminae were examined with an isotope ratio mass spectrometer (DELTA V Plus, Thermo Fisher Scientific Inc., Cambridge, UK). More detailed information is described in the supplementary information.

- Graham, J. B., Dudley, R., Aguilar, N. & Gans, C. Implications of the late Palaeozoic oxygen pulse for physiology and evolution. *Nature* **375**, 117–120 (1995).

2. Dudley, R. Atmospheric oxygen, giant Paleozoic insects and the evolution of aerial locomotor performance. *J. Exp. Biol.* **201**, 1043–1050 (1998).
3. Ainsworth, E. A. & Rogers A. The response of photosynthesis and stomatal conductance to rising [CO₂]: mechanisms and environmental interactions. *Plant, Cell and Environ.* **30**, 258–270 (2007).
4. Tipple, B. J. & Pagani, M. The early origins of terrestrial C₄ photosynthesis. *Annu. Rev. Earth Planet. Sci.* **35**, 435–461 (2007).
5. Stitt, M. Rising carbon dioxide levels and their potential significance for carbon flow in photosynthetic cells. *Plant Cell. Environ.* **14**, 741–762 (1991).
6. Björkman, O. Responses to different quantum flux densities. In *Encyclopedia of Plant Physiology Vol. 12A* (eds Lange, O. L., Nobel, P. S., Osmond, C. B. & Ziegler, H.) 277–338 (Springer, Heidelberg, 1981).
7. Hatch, M. D. C₄ photosynthesis—a unique blend of modified biochemistry, anatomy and ultrastructure. *Biochim. Biophys. Acta.* **895**, 81–106 (1987).
8. Ehleringer, J. R., Cerling, T. E. & Helliker, B. R. C₄ photosynthesis, atmospheric CO₂, and climate. *Oecologia* **112**, 285–299 (1997).
9. Osbome, C. P. & Sack L. Evolution of C₄ plants: a new hypothesis for an interaction of CO₂ and water relations mediated by plant hydraulics. *Phil. Trans R. Soc. B* **367**, 583–600 (2012).
10. Brooks, J. R., Flanagan, L. B., Varney, G. T. & Ehleringer, J. R. Vertical gradients in photosynthetic gas exchange characteristics and refixation of respired CO₂ within boreal forest canopies. *Tree Physiol.* **17**, 1–12 (1997).
11. Pfanz, H. *et al.* Photosynthetic performance (CO₂-compensation point, carboxylation efficiency, and net photosynthesis) of timothy grass (*Phleum pratense* L.) is affected by elevated carbon dioxide in post-volcanic mofette areas. *Environ. Exp. Bot.* **61**, 41–48 (2007).
12. Pearcy, R. W. Photosynthetic gas exchange responses of Australian tropical forest trees in canopy, gap and understory microenvironments. *Funct. Ecol.* **1**, 169–178 (1987).
13. Chazdon, R. L. Sunflecks and their importance to forest understorey plants. *Adv. Ecol. Res.* **18**, 1–63 (1988).
14. Ishida, A., Toma, T., Mori, S. & Marjenah. Effects of Foliar nitrogen and water deficit on the carbon economy of *Shorea smithiana* Sym. seedlings. *Biotropica* **32**, 351–358 (2000).
15. Sternberg, L. *et al.* Carbon dioxide recycling in two Amazonian tropical forests. *Agric. Forest Meteor.* **88**, 259–268 (1997).
16. Kondo, M., Muraoka, H., Uchida, M., Yazaki, Y. & Koizumi, H. Refixation of respired CO₂ by understory vegetation in a cool-temperate deciduous forest in Japan. *Agric. Forest Meteor.* **134**, 110–121 (2005).
17. Pinker, R. The microclimate of a dry tropical forest. *Agri. Meteor.* **22**, 249–265.
18. Körner, C. H. *Scaling from species to vegetation: The usefulness of functional groups.* In *Biodiversity and Ecosystem Function* (eds Schulze, E. –D. & Mooney, H. A.) 117–140 (Springer, Heidelberg, 1994).
19. Mori, S. *et al.* Mixed-power scaling of whole-plant respiration from seedlings to giant trees. *Proc. Natl. Acad. Sci. USA* **107**, 1447–1451 (2010).
20. Adachi, M., Ishida, A., Bunyavejchewin, S., Okuda, T. & Koizumi, H. Spatial and temporal variation in soil respiration in a seasonal dry tropical forest, Thailand. *J. Trop. Ecol.* **25**, 531–539 (2009).
21. Ishida, A., Nakano, T., Matsumoto, Y., Sakoda, M. & Ang, L. H. Diurnal changes in leaf gas exchange and chlorophyll fluorescence in tropical tree species with contrasting light requirements. *Ecol. Res.* **14**, 77–88 (1999).
22. Yamashita, N., Ishida, A., Kushima, H. & Tanaka, N. Acclimation to sudden increase in light favoring an invasive over native trees in subtropical islands, Japan. *Oecologia* **125**, 412–419 (2000).
23. Araki, M. G. *et al.* Estimation of whole-stem respiration, incorporating vertical and seasonal variations in stem CO₂ efflux rate, of *Chamaecyparis obtusa* trees. *J. For. Res.* **15**, 115–122 (2010).
24. Robertson, A. L. *et al.* Stem respiration in tropical forests along an elevation gradient in the Amazon and Andes. *Global Change Biol.* **16**, 3193–3204 (2010).
25. Zach, A., Horna, V. & Leuschner, C. Diverging temperature response of tree stem CO₂ release under dry and wet season conditions in a tropical montane moist forest. *Trees* **24**, 285–296 (2010).
26. De Schepper, V., De Swaef, T., Bauweraerts, I. & Steppe, K. Phloem transport: a review of mechanisms and controls. *J. Exp. Bot.* **64**, 4839–4850 (2013).
27. Negishi, K. Bark respiration rate in stem segments detached from young *Pinus densiflora* trees in relation to velocity of artificial sap flow. *J. Jpn. For. Soc.* **61**, 88–93 (1979).
28. Teskey, R. O., Saveyn, A., Steppe, K. & McGuire, M. A. Origin, fate and significance of CO₂ in tree stems. *New Phytol.* **177**, 17–32. (2008).
29. Bekku, S. Y., Sakata, T., Tanaka, T. & Nakano, T. Midday depression of tree root respiration in relation to leaf transpiration. *Ecol. Res.* **26**, 791–799 (2011).
30. Kuzyakov, Y. & Gavrichkova O. Time lag between photosynthesis and carbon dioxide efflux from soil: a review of mechanisms and controls. *Global Change Biol.* **16**, 3386–3406 (2010).
31. Hanba, Y., Mori, S., Lei, T. T., Koike, T. & Wada, E. Variations in leaf δ¹³C along a vertical profile of irradiance in a temperate Japanese forest. *Oecologia* **110**, 253–261 (1997).
32. Ishida, A. *et al.* Photoprotection of evergreen and drought-deciduous tree leaves to overcome the dry season in monsoonal tropical dry forests in Thailand. *Tree Physiol.* **34**, 15–28 (2014).

Acknowledgments

This study was supported by grants-in-aid from the Japan Society for the Promotion of Science (nos. 18255011, 24370009 to A.I.; 22255004, 22370010 to J.Y.). We wish to thank the staff of the research station, Dr. T. Artchawakom, and Dr. M. Takahashi for their support.

Author contributions

A.I. and T.N. designed and carried out the major part of the field measurements. M.A., K.Y., N.O. and P.L. carried out the field measurements. S.D. and L.P. designed and prepared the field works. A.I. and J.Y. wrote the manuscript.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Ishida, A. *et al.* Effective use of high CO₂ efflux at the soil surface in a tropical understory plant. *Sci. Rep.* **5**, 8991; DOI:10.1038/srep08991 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

Supplementary Information

Title: Effective use of high CO₂ efflux at soil surface in a tropical understory plant

Authors: Atsushi Ishida, Takashi Nakano, Minaco Adachi, Kenichi Yoshimura, Noriyuki Osada, Phanumard Ladpala, Sapit Diloksumpun, Ladawan Puangchit, Jin Yoshimura

Environmental description and more detailed methods

Study site and plant materials

The study was located in the dry evergreen forest at the Sakaerat Environmental Research Station (14° 29'N, 101° 55'E, 563 m ASL), approximately 180 km northeast of Bangkok in Thailand. Mean annual temperature was 26.2°C, and mean annual rainfall was 1240 mm (Sakurai *et al.* 1998). There is a distinct dry season from November to March (Ishida *et al.* 2006). The soil is of sandstone origin and acidic (around 4.5 in pH), and has relatively poor nutrients with high porosity. *Hopea ferrea* Lanessan (Dipterocarpaceae) is the predominant tree with tall canopies (approximately 25-35 m high) in the evergreen forest. Details in landform and soil characteristics are shown in Pitman (1996) and Murata *et al.* (2009).

The examined ginger plant (*Kaempferia marginata* Carey, Zingiberaceae) is a drought-deciduous perennial herb. The plants are usually found near the roadside in the dry evergreen forests with dense canopies, and inside the drought deciduous forests with sparse canopies in the Sakaerat Environmental Research Station. Thus, the ginger plant seems to favor to a relatively light understory.

Measurements of leaf size distribution

We selected a population of the ginger plant found roadside in the dry evergreen forest. To examine the ontogenetic variations of leaf form, we measured the leaf area, leaf length and width, and the top heights of the leaf blades in 150 individual plants with contrasting plants size. The data in leaf shape and the top height are shown in Supplementary Fig. 1 and Supplementary Table 1, respectively.

Microclimate measurements

Photosynthetic photon flux (PPF) was measured with quantum sensors (LI-190SB, LI-COR, Lincoln, NE, USA) at an open place and two understory places around the center of a ginger plant population for three consecutive, relatively sunny days from July 15 to July 17 in 2008 in the middle of the wet season. PPF measurement at the open site was conducted on the top of a 45-m high scaffolding tower constructed near the study site, which exceeded the uppermost canopy in the forest. Ambient air temperature (T_{air}) and relative humidity (RH) in the understory were simultaneously measured with thermistor and thin-film capacitance sensors, respectively (Model 36355, Hioki-Denki, Nagano, Japan). These sensors were connected with small data loggers (Model 3631 or 3635, Hioki-Denki), and the data were minutely stored. These measurements in the understory were conducted at 20 cm above the ground near the center of the ginger population.

On a sunny day (July 15 in 2008), the diurnal variations in ambient air CO₂ concentrations ([CO₂]) were directly measured with thin-film capacitance CO₂ sensors (GM70, Vaisala, Helsinki, Finland) without tube-absorbing air. The CO₂ sensors were set at two heights: (1) 20 cm above the ground and (2) in the air space between the leaf blade and the ground surface in a lamina. To measure CO₂ concentrations in the air space, we selected an individual plant with a relatively large leaf area to avoid air leak along the side of the sensor probe (Supplementary Figure 4).

Environmental description

Supplementary Figure 2 shows the diurnal time variations in microclimate (PPF, air temperature, and relative humidity in air) in the examined population growing at a roadside in the dry evergreen forest and the diurnal variations in PPF at an open place. The daily total PPF at the study site relative to that of the open site was 6.4%. The understory light levels in tropical evergreen forest are approximately 1% of full sunlight or sometimes less than 1% (e.g., Ashton 1992). Because the ginger plants are not found in deeper-shaded sites in the evergreen forests, the light levels of approximately 6.4% relative to full sunlight appear to be required to maintain a population of the ginger plant.

More detailed methods in the photosynthetic capacity measurements

The leaf photosynthetic capacity in eight individuals was measured with an open, portable measurement system (LI-6400, LI-COR, Lincoln, NE). The measurement was conducted in six healthy, mature leaves. The leaf chamber with 6 cm² was used and the red-blue RED lamp unit was utilized as a light source. Photosynthetic light-response curves were measured on the same eight leaf blades under 600 and 400 μmol mol⁻¹ CO₂ in the inlet gas stream with LI-6400. The values of 400 and 600 μmol mol⁻¹ CO₂ were approximate and corresponded to daily mean air CO₂ concentrations at the 20 cm above and just below the leaf blades, respectively. PPFs were decreased stepwise from 800, 500, 200, 100, 70, 40, 30, 20, 10, 7, 3, to 0 μmol m⁻² s⁻¹. Light compensation points and apparent quantum use efficiencies were calculated from linear regressions under very low PPFs (from 0 to 10 μmol m⁻² s⁻¹). The mean leaf temperature during these measurements was 29.2°C which approximately corresponds to the daytime leaf temperature. Photosynthetic ambient air CO₂-response curves were measured on the same seven leaf blades under 500 and 40 μmol m⁻² s⁻¹ PPF with LI-6400. In the values that were exposed to sun-flecks and in those without sunflecks during the daytime, the values of 500 and 40 μmol m⁻² s⁻¹ PPF were used, respectively (see Supplementary Fig. 2). The [CO₂] in the inlet gas stream with LI-6400 increased stepwise from 0, 50, 100, 200, 300, 400, 500, 600, 700, 800, to 1000 μmol mol⁻¹.

More detailed methods in the diurnal variations of leaf gas exchange and chlorophyll fluorescence measurements

The diurnal time change in leaf gas exchange was measured with an open, portable measurement system (LI-6400, LI-COR, Lincoln, NE), from dawn to dusk on 15 July 2008. The measurement was conducted in eight healthy, mature leaves at approximately 30-minute intervals. The leaf chamber with 6 cm² was used and the top part of the chamber was sealed with a clear plastic plate to receive naturally incident PPF. The CO₂ concentration in the inlet gas stream within LI-6400 was adjusted at

600 $\mu\text{mol mol}^{-1}$, which approximately corresponded to the mean air $[\text{CO}_2]$ just below the leaf blades (see Figure 2).

While measuring the leaf gas exchange, the diurnal time variations in chlorophyll *a* fluorescence were measured with a fluorescence meter (Mini-PAM, Walz, Effeltrich, Germany), according to Bilger *et al.* (1995). The fiber-optic cable was connected with the clear top-cover of the LI-6400 chamber, while the angle (60°) and the distance between the leaf surface and the fiber-optic cable were manually adjusted. Maximum fluorescence yield (F_m) and dark fluorescence yield (F_o) in photosystem II (PSII) were determined just before dawn. Just after the measurement of leaf gas exchange, we supplied a saturated-light pulse to the leaf surface. Maximum fluorescence (F_m') and steady-state fluorescence (F) in the light-adapted state of PSII were measured during the daytime. Chlorophyll fluorescence parameters were calculated, according to Genty *et al.* (1989). The potential maximum quantum yield of PSII ($F_v/F_m = (F_m - F_o)/F_m$) was calculated from the dark-time measurements made before dawn. For each daytime measurement, the effective quantum yield of PSII ($\Phi_{\text{PSII}} = (F_m' - F)/F_m'$) was calculated. Assuming that photosystem I and II absorb equal amounts of light and the leaf absorbance of lamina is 0.84, the electron transport rate through PSII (ETR) was calculated as, $\text{ETR} = 0.5 \Phi_{\text{PSII}} 0.84 \text{ PPF}$ (at the leaf surface). Non-photochemical quenching ($\text{NPQ} = (F_m/F_m') - 1$) was also calculated. Data in chlorophyll fluorescence are showed in Supplementary Figure 3.

Measurements of the nitrogen and stable carbon isotope ratio in lamina and the number and size of stomata


After all measurements, we collected the leaves and then cut leaf discs with a borer. The leaf discs were oven dried (70°C , 72 hr) and weighed to determine leaf dry mass per unit leaf area (LMA). The total nitrogen (N) and carbon (C) contents within the leaf discs were measured with an N-C analyzer (Sumigraph NC-900, Sumitomo-Kagaku, Osaka).

To estimate the averaged internal CO_2 concentrations in leaves for a long time, the stable carbon isotope ratios ($\delta^{13}\text{C}$) in lamina were determined with an isotope ratio mass spectrometer (DELTA V Plus, Thermo Fisher Scientific Inc., Cambridge, UK). The $\delta^{13}\text{C}$ values were expressed in delta notation relative to a PD Belemnite standard: $\delta^{13}\text{C} = (R_{\text{sample}} - R_{\text{standard}} / R_{\text{standard}}) \times 1000$ (‰), where R_{sample} is the $^{13}\text{C}/^{12}\text{C}$ ratios of the samples and R_{standard} is the $^{13}\text{C}/^{12}\text{C}$ ratio of the standard.


The numbers and the pore length of stomata in the adaxial and abaxial leaf surfaces were determined by obtaining replicas of the surface of four healthy leaves with a celluloid plate (Universal Micro-printing, SUMP, Tokyo, Japan).

Supplementary Table 1: The morphological and physiological characteristics of leaf blades.


Variables	unit	mean	1 SD
Top height of leaf blades above the ground	mm	24	13
Area of individual leaf blades	cm ²	65.0	36.9
Length/width ratio in individual leaf blades		1.32	0.30
Leaf dry mass per leaf area (LMA)	g m ⁻²	35.8	4.7
Nitrogen (N) content per leaf area	mmol N m ⁻²	54.8	6.4
N concentration per leaf dry mass	mmol N g ⁻¹	1.54	0.15
C/N ratio in leaf blades		19.9	2.05
Stomatal pore length	μm	39.75	6.37
Stomatal density in adaxial leaf surface	No. mm ⁻²	1.6	0.6
Stomatal density in abaxial leaf surface	No. mm ⁻²	21.0	4.3
Light compensation point	μmol m ⁻² s ⁻¹	2.5	1.5
Apparent quantum use efficiency		0.071	0.008
Area-based dark respiration rates	μmol m ⁻² s ⁻¹	-0.17	0.08



Supplementary Figure 1: The ontogenetic variations in size and shape of leaf blades in 150 ginger individuals with different plant sizes. (a) The frequency of blade size of single leaves, and (b) the change of leaf shape (the ratio of length to width in each leaf blade) with leaf area.



Supplementary Figure 2: The diurnal time courses in microclimate at the understory during the successive three days from on 15 July to 17 July (the mid-rainy season) in 2008. Incident photosynthetic photon flux (PPF) at (a) an open place and (b) the understory (20 cm above the ground), and (c) air temperature (T_{air}) and relative humidity (RH) in air at an understory site.



Supplementary Figure 3: The relationships between photosynthetic capacity and photosynthetic photon flux (PPF) at the leaf surface. Data were obtained from the measurements of diurnal time courses in (a) net photosynthetic rates, (b) PSII quantum yield, (c) electron transport rates through PSII, and (d) Stern-Volmer non-photochemical quenching coefficient.



Supplementary Figure 4: The measurement of CO₂ concentrations in the air space between the leaf blade and the soil surface.

References

- Ashton, P. M. S. Some measurements of the microclimate within a Sri Lankan tropical rainforest. *Agri. Forest Meteorol.* **59**, 217-235 (1992).
- Bilger, W., Schreiber, U. & Bock, M. Determination of the quantum efficiency of photosystem II and of non-photochemical quenching of chlorophyll fluorescence in the field. *Oecologia* **102**, 425-432 (1995).
- Genty, B., Briantais, J. -M. & Baker, N. R. The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochem. Biophys. Acta* **990**, 87-92 (1989).
- Ishida, A., Diloksumpun, S., Ladpala, P., Staporn, D., Panuthai, S., Gamo, M., Yazaki, K., Ishizuka, M. & Puangchit, L. Contrasting seasonal leaf habits of canopy trees between tropical dry-deciduous and evergreen forests in Thailand. *Tree Physiol.* **26**, 643-656 (2006).
- Murata, N., Ohta, S., Ishida, A., Kanzaki, M., Wachirinrat, C., Archawakom, T. & Sase, H. Comparison of soil depths between evergreen and deciduous forests as a determinant of their distribution, Northeast Thailand. *J. Forest Res.* **14**, 212-220 (2009).
- Pitman, J. I. Ecophysiology of tropical dry evergreen forest, Thailand: measured and modelled stomatal conductance of *Hopea ferrea*, a dominant canopy emergent. *J. Appl. Ecol.* **33**, 1366-1378 (1996).
- Sakurai, K., Tanaka, S., Ishizuka, S., Kanzaki, M. Differences in soil properties of dry evergreen and dry deciduous forests in the Sakaerat Environmental Research Station. *Tropics* **8**, 61-80 (1998).